

BODY PEEL

Glycolic Acid Gel 70%

INDICATION

Dry, hypotonic or sagging skin, recent or elder stretch marks. Age or seborrhoyc keratosis and ichthyosis.

HOME PRE-TREATMENT

It is advisable to use home treatment products containing glycolic acid for at least 10 to 15 days before the peeling. This will generate a preliminar skin exfoliation, will normalize the skin's pH at physiological values (4-5) and will enable a first evaluation of the skin's tolerance to glycolic acid.

Precautions

Inform patient to avoid using depilatory waxes, electrodepilation and other abrasive or irritating substances during the 2 or 3 days before the peeling.

PRE-PEELING PHASE

Cheratolitic Pre-peeling is a solution with salicylic acid, lactic acid and glycolic acid. Its purpose is to enable the subsequent penetration of the glycolic acid in the following phase. Apply the solution with delicate movements in the stretch-marks or in the interested area do not rinse let it dry.

Precautions

Avoid using the *Cheratolitic Pre-peeling* in particularly sensitive areas or on broken skin. Use with care over large surfaces due to the presence of salicylic acid.

PEELING PHASE

Glycolic Acid Gel 70% free at pH 1,6. Chemical agent capable of stimulating the fibroblast to produce new collagen in the derm. Its presentation as a dense gel allows a handy application. Apply with a glove, with a short massage to obtain uniformity and penetration. The purpose is to obtain a pink uniform erithema.

NEUTRALIZING PHASE

The acid's activity is stopped by rinsing thoroughly with abundant fresh water. Neutralizer is also useful at any time during the peeling phase whenever a premature penetration of the acid occurs in a specific area.

POST-PEELING PHASE

The specific product *Stand-by Cream* is indicated to heal and nourish the stimulated or irritated skin after a peeling. Apply 2 or 3 times a day for 3 or 4 days. *Stand-by Cream* can be used for every further dry skin symptoms.

Precautions

Inform the patient to avoid scratching, pulling or submit to irritating stimulations the treated area. The use of home care treatment products should start again after 3 to 5 days. The use of *Advanced Lotion* with glycolic acid at 30% can be recommended under medical supervision.

PEELING SEQUENCE

It is advisable to repeat the peeling every 12 to 15 days.